

### **REMARKS**

The present application relates to inbred maize plant and seed PH0GC. Claims 1-30 are pending in the present application. No new matter has been added by way of amendment. Applicant respectfully requests consideration of the claims in view of the following remarks.

#### **Detailed Action**

Applicant acknowledges that because this application is eligible for continued examination under 37 C.F.R. § 1.114 and the fee set forth in 37 C.F.R. § 1.17(e) has been timely paid, the finality of the previous Office Action has been withdrawn pursuant to 37 C.F.R. § 1.114. Applicant further acknowledges that Applicant's submission filed on October 21, 2005 has been entered.

#### **Request for Information under 37 C.F.R. § 1.105**

The Examiner has made a Request for Information under 37 C.F.R. § 1.105. The Examiner states the requested information is "required to make a meaningful and complete search of the prior art". See Office Action, pp. 2-3 and 13-15.

Applicant provides answers to each of the Examiner's interrogatories discussed *infra*. Applicant notes that the information provided to the third and fourth interrogatories are only to *previously* publicly disclosed or sold parental maize lines or progeny therefrom as requested by the Examiner. Thus, Applicant asserts the interrogatories have been answered with respect to the Examiner's request for the information for prior art purposes. Applicant points out that the third interrogatory was specific to *previously* publicly disclosed or sold as this is relevant to the Examiner's prior art inquiry. Thus Applicant notes that the response to the fourth interrogatory is also answered with respect to maize lines produced by said method using said original parental maize lines which were *previously* publicly disclosed, sold or disclosed in a U.S. patent application as this is relevant to the Examiner's request for prior art purposes as stated on page 13 of the Office Action.

The Examiner begins by asking firstly, what were the original parental maize lines used to produce maize inbred line PH0GC? PH0GC was derived from a synthetic population named SYN92F.

Secondly, what method and steps were used to produce maize inbred line PH0GC? Pedigree selection method produced from SYN92F by selfing and ear rowing from F0 through F11 generation.

Third, have any of said parental maize lines (a) or progeny (b) therefrom been previously publicly disclosed or sold?

a. Pioneer Hi-Bred has not previously publicly disclosed or sold the synthetic SYN92F.

b. Pioneer Hi-Bred has not previously publicly disclosed or sold progeny of the synthetic SYN92F prior to the earliest priority date.

Fourth, were any other maize lines produced by said method using said original parental maize lines, and if so, have said produced maize lines been publicly disclosed, sold or disclosed in a U.S. patent application? If so, under what designation were said other maize lines disclosed or sold? No maize line using the synthetic SYN92F has been *previously* publicly disclosed, sold or disclosed in a U.S. patent application by Applicant prior to the earliest priority date.

In light of the above remarks, Applicant respectfully requests reconsideration and compliance with the interrogatories under the Request for Information under 37 C.F.R. § 1.105.

#### **Rejections Under 35 U.S.C. § 112, Second Paragraph**

Claims 11-30 stand rejected under 35 U.S.C. § 112, second paragraph, as being indefinite for failing to particularly point out and distinctly claim the subject matter which Applicant regards as the invention. *See Office Action*, pp. 3-4.

The Examiner states claim 11 is indefinite "because it is unclear what the metes and bounds of a maize plant having all of the physiological and morphological characteristics of inbred maize line PH0GC are."

Applicant traverses this rejection. Claim 11 specifically claims a maize plant having all the physiological and morphological characteristics of inbred line PH0GC. Claim 11 encompasses maize plants having the characteristics of inbred line PH0GC. Applicant believes the Examiner is making the assumption that the fact that one must use seed of the maize inbred line PH0GC itself to obtain a plant with the same morphological and physiological characteristics as a plant of the variety PH0GC. However, one of ordinary skill in the art can obtain a plant with all of the same morphological and physiological characteristics as maize inbred line PH0GC without actually using seed of maize inbred line PH0GC. For example, this

can be accomplished by using double haploid technology to "recreate" PH0GC through the use of F1 hybrid seed in which PH0GC was a parent. As emphasized in previous office action responses, all members of the genus of F1 hybrids seed will receive one non-recombinant set of chromosomes of PH0GC. By using the seed of an F1 hybrid made with PH0GC, one can recover this non-recombined set of chromosomes from the F1 hybrid seed. Thus, a plant that has all of the same morphological and physical characteristics of PH0GC can be created without direct use of seed of inbred line PH0GC. Applicant directs the Examiner to the following web site which further explains and illustrates double haploid technology at the internet address [www.uni-hohenheim.de/%7Eipspwww/350b/indexe.html#Project3](http://www.uni-hohenheim.de/%7Eipspwww/350b/indexe.html#Project3) (attached as Appendix 1), as well as to U.S. Patent No. 5,770,788 to Jia and U.S. Patent No. 6,200,808 to Simmonds *et al.*. As noted on the web site, the use of double haploid technology to has been used in plant breeding to produce desired homozygous inbred lines for more than 50 years.

Claim 25 is rejected as indefinite "because it is directed to a maize plant derived from inbred line PH0GC, but is dependent upon claim 11 that is not specifically directed to inbred line PH0GC, only a maize plant having all of the physiological and morphological characteristics of PH0GC".

Applicant traverses this rejection for the reasons asserted *supra*. Claim 25 is definite and does include the plant of claim 11 wherein the plant is "[a] maize plant having all the physiological and morphological characteristics of inbred line PH0GC, wherein a sample of the seed of inbred line PH0GC was deposited under ATCC Accession Number PTA-4523". In addition, claim 25 claims the maize plant of claim 11 with these additional limitations, which are not necessarily present in the maize plant of claim 11. The presence of these additional limitations does not mean that claim 25 does not possess all limitations of claim 11; these claims still require a maize plant having the physiological and morphological characteristics of inbred line PH0GC. Because claim 25 does incorporate all elements of claim 11, it is in accordance with the requirements of § 112, second paragraph.

Claims 28 and 29 are rejected as indefinite "because it is unclear what the metes and bounds of employing the maize plant of claim 11 are".

Applicant traverses this rejection for the reasons asserted *supra*. Claims 28 and 29 are definite and do include the plant of claim 11. Thus, because claims 28 and 29 do incorporate all elements of claim 11, it is in accordance with the requirements of § 112, second paragraph.

Claim 30 is rejected as the Examiner states it is indefinite "because the method requires 'obtaining an F1 hybrid seed from which maize inbred line PH0GC is a parent', but said claim is dependent upon claim 11 which is directed to a 'maize plant having all the physiological and morphological characteristics of inbred line PH0GC'".

Applicant traverses this rejection for the reasons asserted *supra*. Claim 30 is definite and does incorporate all elements of claim 11, and therefore it is in accordance with the requirements of § 112, second paragraph.

In light of the above amendments and remarks, Applicant respectfully requests reconsideration and withdrawal of the rejections under 35 U.S.C. § 112, second paragraph.

#### **Rejections Under 35 U.S.C. § 112, First Paragraph**

##### *A. Written description regarding Claims 1-10 and 11-30*

Claims 1-10 remain rejected and claims 11-30 stand rejected under 35 U.S.C. § 112, first paragraph, as failing to comply with the written description requirement. The claims(s) contains subject matter, which was not described in the specification in such a way as reasonably convey to one skilled in the relevant art that the inventor(s), at the time the application was filed, had possession of the claimed invention. The Examiner states the rejection is repeated for the reasons of record set forth in the Office Action of July 21, 2005. *See* Office Action, pp. 4-8.

Applicant respectfully traverses this rejection. Applicant reiterates that the written description requirement of § 112, first paragraph has been fulfilled by depositing seeds of PH0GC in a public depository and by referencing the deposit in the specification. *See* specification, p. 76, ll. 2-28; *see also Enzo Biochem, Inc. v. Gen-Probe Inc.*, 323 F.3d 956, 965, 63 U.S.P.Q.2d 1609, 1613 (Fed. Cir. 2002) (stating that the written description requirement of § 112, ¶ 1 may be fulfilled by depositing material in a public depository, where the deposited material is not accessible in writing, and where reference to the deposit is made in the specification). This deposit not only describes inbred maize line PH0GC but also the hybrid maize plants, plant parts, and seeds grown in claims 1-10 and 11-30. In a prior case before the Board of Patent Appeals and Interferences, the Board determined that where claims to an inbred maize plant satisfied the written description requirement, claims to the F1 hybrid seed and plants with the inbred maize plant as a parent also satisfied the written description requirement. *See Ex parte Carlson* (B.P.A.I. 2005). The Board therein stated:

All that is required by the claims is that the hybrid has one parent that is a plant of corn variety [inbred]. Since the examiner has indicated that the seed and the plant of the corn variety [inbred] are allowable . . . there can be no doubt that the specification provides and adequate written description of this corn variety. In addition, the examiner appears to recognize (Answer, page 25) that appellant's specification describes an exemplary hybrid wherein one parent was a plant of the corn variety [inbred]. . . Accordingly, it is unclear to this merits panel what additional description is necessary.

*Ex parte Carlson*, p. 16. Here, Applicant has done just what the applicant in *Ex parte Carlson* did, that is claim hybrids having one parent that is a plant of an inbred variety. Further, Applicant reiterates that the specification contains an example of a hybrid produced by PH0GC in the application as filed. See specification, p. 38, Table 3. Thus, under *Ex parte Carlson*, "it is unclear . . . what additional description is necessary." See *Ex parte Carlson*, p. 16; see also *Regents of Univ. of Cal.*, 119 F.3d at 1569, 43 U.S.P.Q.2d at 1406 (stating that an Applicant is "not required to disclose every species encompassed by their claims even in an unpredictable art").

Applicant reiterates that each member of the genus of hybrids which has PH0GC has a parent and which is encompassed by claims 1-10 and 11-30 shares the identifying structural feature of the cells and/or chromosomes of inbred line PH0GC. An Applicant's claims are described where they set forth and define "structural features commonly possessed by members of the genus that distinguish them from others." *Regents of Univ. of Cal. v. Eli Lilly & Co.*, 119 F.3d 1559, 1568, 43 U.S.P.Q.2d 1398, 1406 (Fed. Cir. 1997) (emphasis added). One of skill in the art, utilizing technology well known in the art, could identify any member of the claimed genus.

The Examiner again cites *In re Wallach*, 71 USPQ2d 1939 at 1940 (C.A.F.C. 2004).

Applicant respectfully traverses and reiterates that the *Wallach* case is not applicable to the claimed invention. Unlike in *Wallach*, the issue in the present case is the characterization of an entire genome, not a single isolated protein. Those of skill in the art utilize molecular markers, such as SSR's, to characterize plant genomes. As Applicant clearly teaches in the specification:

"In addition to phenotypic observations, a plant can also be identified by its genotype. The genotype of a plant can be characterized through a genetic marker profile, which can identify plants of the same variety or a related variety or be used to determine or validate a pedigree. Genetic marker profiles can be obtained by techniques such as... Simple Sequence Repeats (SSRs)...For

example, see Berry, Don, et al., "Assessing Probability of Ancestry Using Simple Sequence Repeat Profiles: Applications to Maize Hybrids and Inbreds", *Genetics*, 2002, 161:813-824, which is incorporated by reference herein." See specification, p. 16, lines 23.

The use of molecular marker profiles by those of ordinary skill in the art in backcrossing is also clearly supported by the scientific literature. For example, see Ragot, M. *et al.* (1995) Marker-assisted backcrossing: a practical example, in *Techniques et Utilisations des Marqueurs Moléculaires (Les Colloques*, Vol. 72, pp. 45-56 (attached as Appendix 2), and Openshaw *et al.*, (1994) Marker-assisted Selection in Backcross Breeding, Analysis of Molecular Marker Data, pp. 41-43 (attached as Appendix 3). Specifically, Ragot *et al.* concludes that "recovery of the recurrent parent genotype could proceed even faster than in the experiment described herein, should the appropriate protocol and resources (population size, number and position of markers) be allocated." Therefore, one of ordinary skill in the art can obtain the unique SSR profile of PH0GC which can be used to identify essentially derived varieties and other progeny lines developed from the use of PH0GC, as well as cells and other plant parts thereof.

The Examiner further states that the instant disclosure "only provides an adequate written description for inbred maize line PH0GC, because the functional characteristics of an F1 progeny would be correlated as much with the second parent as with inbred maize line PH0GC". See Office Action, pp. 6-7.

Applicant respectfully traverses this rejection. Applicant reiterates that each F1 hybrid which has PH0GC as a parent and which is encompassed by claims 1-10 and 11-30 contain at least one set of chromosomes of inbred line PH0GC. Thus, these claims set forth "structural features commonly possessed by members of the genus that distinguish them from others," as only F1 hybrids with PH0GC as a parent would have a complete set of PH0GC chromosomes. *Regents of Univ. of Cal.*, 119 F.3d at 1568, 43 U.S.P.Q.2d at 1406. The claimed F1 hybrids are therefore described in such a way that distinguishes them from other hybrids, which is sufficient to meet the written description requirement. See *id.*

Further, at its foundation, the written description requirement serves an evidentiary function of making certain that the Applicant is in possession of a specific characteristic that identifies their claimed invention. The data provided by Applicant in Tables 1 and 2-4 serves this purpose. See specification, pp. 18-20 Table 1; pp. 37-39 Tables 2-4. The other inbred is not

the point of patentability, nor is it what is being claimed. Rather, the relevant claims are drawn precisely to what is described, inbred maize line PH0GC including the hybrid maize plants, plant parts, and seeds grown in claims 1-10 and 11-30.

It is undisputed that fingerprinting with molecular markers is widely used for characterizing germplasm. Specifically, SSR profiles are known and can be practiced by one of ordinary skill in the art in maize breeding. One of ordinary skill has been enabled by the deposit to make and use minor variants of inbred maize line PH0GC, and one of ordinary skill in the art uses SSR markers to characterize backcross conversions of an inbred. Applicant has claimed in the manner used by those of ordinary skill in the art to characterize backcross conversions. Thus, Applicant respectfully submits the claimed invention is in accordance with the written description guidelines.

One skilled in the art would thus recognize that Applicant was in possession of the invention described in claims 1-10 and 11-30 as of the filing date of the application. Accordingly, Applicant respectfully requests reconsideration and withdrawal of the rejections under 35 U.S.C. § 112, first paragraph.

*B. Enablement regarding Claims 1-10 and claims 11-30*

Claims 1-10 remain rejected and claims 11-30 stand rejected under 35 U.S.C. § 112, first paragraph, as failing to comply with the enablement requirement. The Examiner asserts that the claim(s) contains subject matter, which was not described in the specification in such a way as to enable one skilled in the art to which it pertains, or with which it is most nearly connected, to make and/or use the invention. The Examiner states the "while being enabling for inbred maize line PH0GC, deposited under ATCC Accession No. PTA-4523 and methods of using, does not reasonably provide enablement for a seed comprising at least one set of chromosomes of maize inbred line PH0GC as broadly claimed." The Examiner states the rejection is repeated for the reasons of record set forth in the Office Action of July 21, 2005. *See* Office Action, pp. 8-11.

Applicant respectfully traverses. Applicant maintains the arguments submitted in the previous Amendment of May 17, 2005 regarding the references (Kevern, Carlone, Segebart '719 and Segebart '109) mentioned by the Examiner.

The Applicant further asserts the specification provides a description of how to backcross traits into PH0GC (Specification, p. 22, l. 33 through p. 23, l. 17) and it is understood by those of

skill in the art that backcross conversions are routinely produced and do not represent a substantial change to a variety. The World Seed Organization, on its web site, writes, "[t]he concept of an essentially derived variety was introduced into the 1991 Act of the UPOV Convention in order to avoid plagiarism through mutation, multiple back-crossing and to fill the gap between Plant Breeder's Rights and patents." ASSINSEL, an International breeders association, has published a position paper that refers to a conversion produced by repeated backcrossing of parental lines of hybrid varieties as a "cosmetic modification". As determined by the UPOV Convention, "essentially derived varieties may be obtained for example by the selection of a natural or induced mutant, or of a somaclonal variant, the selection of a variant individual from plants of the initial variety, backcrossing, or transformation by genetic engineering" (emphasis added). Copies of web pages with these quotes are provided in Appendix 4. Thus, it is clear that there is worldwide agreement that by obtaining the seed of a newly developed variety such as PH0GC, and by using such seed for repeated backcrossing in accordance with the current claims, one is producing only a cosmetic modification and plagiarizing the work of the inbred inventor.

The ability of one of ordinary skill in the art to effectively use backcrossing to introgress a single locus conversion is also clearly supported by the scientific literature. For example, see Ragot, M. *et al.* (1995) Marker-assisted backcrossing: a practical example, in *Techniques et Utilisations des Marqueurs Moleculaires (Les Colloques*, Vol. 72, pp. 45-56 (attached as Appendix 2), and Openshaw *et al.*, (1994) Marker-assisted Selection in Backcross Breeding, Analysis of Molecular Marker Data, pp. 41-43 (attached as Appendix 3). Specifically, Ragot *et al.*, demonstrates that "spectacular" progress toward the recurrent parent genotype was obtained with 61 RFLP markers. Ragot *et al.* concludes that "recovery of the recurrent parent genotype could proceed even faster than in the experiment described herein, should the appropriate protocol and resources (population size, number and position of markers) be allocated."

Furthermore, the specification teaches multiple ways of introgressing or transforming a maize plant with various genes which encode specific protein products which confer advantageous traits desired in the plant. (See generally, specification, p. 22-33).

Accordingly, Applicant submits that claims 1-10 and 11-30 are fully enabled and have fully satisfied the legal standards for enablement. Applicant respectfully requests



reconsideration and withdrawal of the enablement rejections under 35 U.S.C. § 112, first paragraph.


### **Conclusion**

In conclusion, Applicant submits in light of the above amendments and remarks, the claims as amended are in a condition for allowance, and reconsideration is respectfully requested. If it is felt that it would aid in prosecution, the Examiner is invited to contact the undersigned at the number indicated to discuss any outstanding issues.

This is a request under the provision of 37 CFR § 1.136(a) to extend the period for filing a response in the above-identified application for two months from April 12, 2006 to June 12, 2006. A request for an extension of time from April 12, 2006 to May 12, 2006 was previously filed with the Amendment filed May 12, 2006. Applicant is a large entity; therefore, please charge Deposit Account Number 26-0084 in the amount of \$330.00 for a 2-month extension of time minus the one month previously paid. No other fees or extensions of time are believed to be due in connection with this amendment; however, consider this a request for any extension inadvertently omitted, and charge any additional fees to Deposit Account No. 26-0084.

Reconsideration and allowance is respectfully requested.

Respectfully submitted,



LILA A. T. AKRAD, Reg. No. 52,550  
McKEE, VOORHEES & SEASE, P.L.C.  
801 Grand Avenue, Suite 3200  
Des Moines, Iowa 50309-2721  
Phone No: (515) 288-3667  
Fax No: (515) 288-1338  
**CUSTOMER NO: 27142**

-LATA/pw-

Attorneys of Record

**Application of the *in-vivo*-haploid induction in hybrid maize breeding**


# 1. Reproductive and genetic investigations on *in-vivo*-haploid induction in maize (*Zea mays* L.)

△ Contact person:  
Prof. Dr. H.H. Geiger ([geigerth@uni-hohenheim.de](mailto:geigerth@uni-hohenheim.de))




DH-Line in generation D<sub>1</sub>

The interest in haploid/double haploid (H/D) techniques has enormously increased in the last years. The introduction of H/DH-techniques in maize breeding programs traces back to the 1950s. Shortly after the first reports of the spontaneous occurrence of H/DH-plants in maize, scientists and breeders started to discuss the application of such homozygous plants in breeding programs and their commercial use. By means of the development of inducers and a method for artificial doubling of chromosome set, the H/DH-technique has been developed in the past years until such an extent that it is being used as a matter of routine by many breeders.



After pollination with an inducer plant, kernels with H-embryo of maternal origin with triploid endosperm arise, together with regularly

double fertilized kernels. Chromosome elimination and parthenogenesis are considered to be the possible biological mechanisms responsible for the occurrence of H-plants. However, chromosome elimination and parthenogenesis exclude each other per definition. Therefore, we chose the neutral term *in-vivo-haploid induction* for the phenomenon mentioned.



Inductor RWS

The aim of our work was to develop a novel inducer line with an increased induction rate. The inducer line **RWS** developed, displays both advantage of a high induction rate and combination of two dominant identification markers: a red stem, and an embryo and endosperm coloration. Inducer **RWS** enables the breeder to use *in-vivo-haploid induction* as an effective tool for development of H/DH-plants with almost any genetic background. The method is less effective with donor genotypes, carrying the above mentioned identification markers or anthozyne inhibitor-genes themselves.

The spontaneous doubling rate in maize ranges from 1-10 %. Therefore an artificial chromosome doubling method to increase the number of fertile DH-plants is essential. The artificial chromosome doubling method, using colchicine as doubling agent, facilitates an effective development of DH lines.



Identification of H/DH-plants based on lacking stem-coloration



H/DH-field

in *Arabidopsis*. In *Methods in Arabidopsis*  
2.

J.M., GOODMAN H.M., KOORNEEF  
NEIDERWITZ E.M., 1993. An integrated  
J., 3, 745-754.

CAROTTE M., MORISAN A., JOUJON  
DUR D., GERAUDAT J., GUILLEY F.,  
OHS R., ORIELLET P., DELBANY M.,  
LICK J., PHILIPPS G., AXELSON M.,  
An inventory of 1152 expressed sequence  
sequences. *Plant J.*, 4 (8), 1051-1059.

SCHMIDT R., CNOFFS G., DHAN C.,  
JANOFF L., SOMMERVILLE C., 1991.  
that of the *Arabidopsis* genome. *Plant J.*,

Mapping RFLP and phenotypic markers in

DOE W.D.S., HANCOCK E.M., GOODMAN  
Map of *Arabidopsis thaliana*. *Plant Cell*,

, 9, 111-127.

Location of an overlapping YAC library of  
340-351.

## Marker-assisted backcrossing: a practical example

M. RAGOT<sup>1</sup>, M. BIASIOLLI<sup>1</sup>, M.F. DELBUT<sup>2</sup>, A. DELL'ORO<sup>1</sup>, L. MALGARINI<sup>1</sup>,  
P. THEVENIN<sup>1</sup>, J. VERNON<sup>2</sup>, J. VIVANT<sup>2</sup>, R. ZIMMERMANN<sup>1</sup> and G. GAY<sup>1</sup>.

<sup>1</sup> Ciba Seeds, CH-4002 Basle, Switzerland

<sup>2</sup> Département d'Amélioration des Plantes, INRA-Domaine d'Epoisses,  
F-21110 Genlis, France

### Summary

That molecular markers allow fast recovery of recurrent parent genotype in backcross programs is well-known. Restriction Fragment Length Polymorphism (RFLP's) were used in maize to introgress by backcross a transgene construct, containing phosphinothricin resistance and insecticidal protein genes, from a transformed parent into an elite inbred line. At each generation plants carrying the transgene construct were selected based on their phosphinothricin resistance, and further characterized with RFLP's. Both maximum recovery of recurrent parent genotype and minimum linkage drag were taken into account for marker-based selection. Embryo rescue was used to shorten generation time. Progress towards recurrent parent genotype was spectacular. Levels of recurrent parent genotype recovery which would normally be observed, in the absence of selection, in the BC<sub>4</sub> generation were obtained at the BC<sub>3</sub> generation, about one year after BC<sub>1</sub> seeds had been planted. Besides the evidence already provided by RFLP's, phenotypic evaluation of the backcross-derived near-isogenic lines will constitute an additional check of the completeness of the conversion.

### Introduction

Backcrossing has been a common breeding practice for as long as elite germplasm has been available. It has mainly been used to introgress single Mendelian traits, such as disease resistance or quality factors, into elite germplasm (Allard 1960; Hallauer and Miranda 1961). One of the most attractive attributes of backcrossing is that it allows to perform targeted modifications without disrupting the existing overall genetic balance of the recurrent parent.

However, production of fully converted near isogenic lines through classical backcrossing procedures is a lengthy procedure. If at all possible. Theoretically, a minimum

of seven classical backcross generations are required to recover more than 99% of recurrent parent genotype, assuming no linkage drag. The attractiveness of classical backcross procedures is therefore substantially diminished for crops, such as maize (*Zea mays* L.), where the turn-over of elite cultivars is very fast. In addition, full recovery of recurrent parent genotype is usually not achieved through classical backcrossing, which may result in deleterious agronomic effects. Murray *et al.* (1988) reported about 90% recurrent parent genotype recovery in two BC<sub>30</sub>-equivalent conversions (A632Rk and A632Rp) of the maize line A632. The conversions had retained respectively 4 and 7 donor fragments in addition to the one carrying the gene of interest.

Reduction in the number of backcross generations needed to obtain fully converted individuals has been shown theoretically, or from simulations, to be achievable through the use of molecular markers (Thakale *et al.* 1989; Hospital *et al.* 1992; Jarosz *et al.* 1994). Because they provide thorough characterization of the genetic variability at each backcross generation, markers allow to take full advantage of this variability by applying the highest possible selection intensity.

Efficiency of marker-assisted backcrossing was investigated through an experiment aimed at introgressing a single genetic factor (a transgene construct) from a donor into a recipient maize line.

## Materials and methods

### Plant Material

A homozygous inbred maize line of Lancaster origin was used as donor parent to introgress its transgene construct, through repeated backcrossing, into a recipient parent from the Stiff Stalk germplasm group. Both parents are proprietary elite lines. The transgene construct carries both a phosphinothricin resistance gene and synthetic genes encoding the exonuclease fragment of the CryIA(b) *Bacillus thuringiensis* protein (Kozick *et al.* 1993). Transformation was achieved through microprojectile bombardment (Kozick *et al.* 1993) and resulted in a single insertion (*Bt locus*), on chromosome 1 (Figure 1).

### Backcross protocol

The F1 progeny of the cross between the donor and the recipient was screened for the presence of the transgene construct by applying Basta, a phosphinothricin-based herbicide, onto each plant. Resistant individuals were then used to generate BC<sub>1</sub> progeny.

For each backcross generation, except the BC<sub>1</sub>, individuals were planted in multipots and sprayed with Basta to eliminate those which did not carry the transgene construct. To avoid the stress resulting from treatment with Basta, BC<sub>1</sub> plants carrying the transgene construct were identified using Southern blots probed with the *par* and *Bt* genes. Resistant plants were transplanted in an open-soil greenhouse and leaf-sampled for molecular marker

analyses. Results of marker analysis flowering. A single plant was rescued and transferred onto the embryo first underwent a greenhouse medium, before being average, four months.

### Molecular marker analysis

Restriction Fragment Length Polymorphism (RFLP) analysis was performed using the same restriction enzymes and probes as in previous studies. Lanes were chosen from among those provided coverage of the entire contained two loci tightly linked recombination units away (Figure BC<sub>1</sub>+1 generation comprised both or tightly linked ones, and additionally selected BC<sub>1</sub> plant was heterozygous independent reference population generation.

### Selection procedure

At each generation plants recurrent-parent-genotype and attempt to integrate both criteria missing values were not included contributed to the selection procedure best ranking one of these for *w* for the BC<sub>1</sub> selection) was available.

## Results and discussion

### Selection for the gene *or*

The observed segregation significantly different ( $P < 0.05$ ).

### Recurrent parent genotype

Statistics for the genotype performed taking the whole genome backcross-derived plant thereof

or more than 99% of recurrent parent of classical backcross such as maize (*Zea mays* L.), on, full recovery of recurrent crossing, which may result in about 90% recurrent parent (Bt and A632Bp) of the maize donor fragments in addition to

ded to obtain fully converted, to be achievable through the (1992; Jarboe et al. 1994). variability at each backcross ability by applying the highest

igned through an experiment construct) from a donor into a

was used as donor parent to using, into a recipient parent proprietary elite lines. The to gene and synthetic genes *harpin* protein (Kozel et al. 1994) and *harpin* (Kozel et al. 1994) (Figure 1).

recipient was screened for the phthalocyanine-based herbicide, as BC<sub>1</sub> progeny. plants were planted in multipots by the transgene construct. To plants carrying the transgene *cp* and *Bt* genes. Resistant sampled for molecular marker

analyses. Results of marker analyses were made available at the latest two weeks after flowering. A single plant was selected, of which all backcross-derived embryos were rescued and transferred onto tissue culture medium. Plantlets that developed from these embryos first underwent a greenhouse acclimation phase, while still growing on tissue culture medium, before being transplanted into multipots. Backcross cycles lasted, on average, four months.

#### Molecular marker analyses

Restriction Fragment Length Polymorphisms (RFLP's) were used to establish genotypes in all four generations. RFLP detection involved either radioactive or chemoluminescent techniques. For the BC<sub>1</sub> generation, 60 marker-enzyme combinations were chosen from among those revealing polymorphism between donor and recipient. They provided coverage of the entire genome, defining intervals of about 25 cM in size, and contained two loci tightly linked to the *Bt* locus, CG320 and CG415, respectively 5 and 16 recombination units away (Figure 1). For subsequent generations, markers analyzed in the BC<sub>n+1</sub> generation comprised both those for which the selected BC<sub>n</sub> plant was heterozygous, or tightly linked ones, and additional ones located in chromosomal segments for which the selected BC<sub>n</sub> plant was heterozygous (Table 1). Marker map positions were obtained from independent reference populations and confirmed by analysis of segregation in the BC<sub>1</sub> generation.

#### Selection procedure

At each generation plants were ranked based both on the percentage of homozygous recurrent-parent-genotype and on the extent of linkage drag around the *Bt* locus, in an attempt to integrate both criteria. Plants for which two or more adjacent markers had missing values were not included in the analyses. Success or failure of the pollinations also contributed to the selection procedure. One single plant was selected at each generation: the best ranking one of those for which a backcross progeny of size 100 or more (50 or more for the BC<sub>3</sub> selection) was available.

### Results and discussion

#### Selection for the gene of interest

The observed segregation ratios for phosphinothricin resistance (Table 1) were not significantly different ( $P < 0.05$ ) from the expected 1:1, as shown by Chi-square tests.

#### Recurrent parent genotype recovery

Statistics for the genotyped plants are summarized in Table 1. Calculations were performed taking the whole genome into account, including the *Bt* locus. The "perfect" backcross-derived plant therefore carries one heterozygous chromosomal segment, that





C0308  
 C0324  
 C0311  
 C0316  
 C0312  
 C0307  
 C0345  
 C0340

■ heterozygous  
 ■■ homozygous recurrent

Figure 1a: Genetic maps of the backcross-derived individuals selected in the first four generations of a marker-assisted backcross program. The locus to be introgressed (B0) is located on chromosome 1.

# SELECTED BC2

1  
 C0308  
 C0324  
 C0311  
 C0316  
 C0312  
 C0307  
 C0345  
 C0340  
 2  
 C0370  
 C0308  
 C0324  
 C0311  
 C0316  
 C0312  
 C0307  
 C0345  
 C0340  
 3  
 C0328  
 C0314  
 C0306  
 C0301  
 C0305  
 C0309  
 C0313  
 C0315  
 C0317  
 C0319  
 C0321  
 C0323  
 C0325  
 C0327  
 C0329  
 C0331  
 C0333  
 C0335  
 C0337  
 C0339  
 C0341  
 C0343  
 C0345  
 C0347  
 C0349  
 C0351  
 C0353  
 C0355  
 C0357  
 C0359  
 C0361  
 C0363  
 C0365  
 C0367  
 C0369  
 C0371  
 C0373  
 C0375  
 C0377  
 C0379  
 C0381  
 C0383  
 C0385  
 C0387  
 C0389  
 C0391  
 C0393  
 C0395  
 C0397  
 C0399  
 C0401  
 C0403  
 C0405  
 C0407  
 C0409  
 C0411  
 C0413  
 C0415  
 C0417  
 C0419  
 C0421  
 C0423  
 C0425  
 C0427  
 C0429  
 C0431  
 C0433  
 C0435  
 C0437  
 C0439  
 C0441  
 C0443  
 C0445  
 C0447  
 C0449  
 C0451  
 C0453  
 C0455  
 C0457  
 C0459  
 C0461  
 C0463  
 C0465  
 C0467  
 C0469  
 C0471  
 C0473  
 C0475  
 C0477  
 C0479  
 C0481  
 C0483  
 C0485  
 C0487  
 C0489  
 C0491  
 C0493  
 C0495  
 C0497  
 C0499  
 C0501  
 C0503  
 C0505  
 C0507  
 C0509  
 C0511  
 C0513  
 C0515  
 C0517  
 C0519  
 C0521  
 C0523  
 C0525  
 C0527  
 C0529  
 C0531  
 C0533  
 C0535  
 C0537  
 C0539  
 C0541  
 C0543  
 C0545  
 C0547  
 C0549  
 C0551  
 C0553  
 C0555  
 C0557  
 C0559  
 C0561  
 C0563  
 C0565  
 C0567  
 C0569  
 C0571  
 C0573  
 C0575  
 C0577  
 C0579  
 C0581  
 C0583  
 C0585  
 C0587  
 C0589  
 C0591  
 C0593  
 C0595  
 C0597  
 C0599  
 C0601  
 C0603  
 C0605  
 C0607  
 C0609  
 C0611  
 C0613  
 C0615  
 C0617  
 C0619  
 C0621  
 C0623  
 C0625  
 C0627  
 C0629  
 C0631  
 C0633  
 C0635  
 C0637  
 C0639  
 C0641  
 C0643  
 C0645  
 C0647  
 C0649  
 C0651  
 C0653  
 C0655  
 C0657  
 C0659  
 C0661  
 C0663  
 C0665  
 C0667  
 C0669  
 C0671  
 C0673  
 C0675  
 C0677  
 C0679  
 C0681  
 C0683  
 C0685  
 C0687  
 C0689  
 C0691  
 C0693  
 C0695  
 C0697  
 C0699  
 C0701  
 C0703  
 C0705  
 C0707  
 C0709  
 C0711  
 C0713  
 C0715  
 C0717  
 C0719  
 C0721  
 C0723  
 C0725  
 C0727  
 C0729  
 C0731  
 C0733  
 C0735  
 C0737  
 C0739  
 C0741  
 C0743  
 C0745  
 C0747  
 C0749  
 C0751  
 C0753  
 C0755  
 C0757  
 C0759  
 C0761  
 C0763  
 C0765  
 C0767  
 C0769  
 C0771  
 C0773  
 C0775  
 C0777  
 C0779  
 C0781  
 C0783  
 C0785  
 C0787  
 C0789  
 C0791  
 C0793  
 C0795  
 C0797  
 C0799  
 C0801  
 C0803  
 C0805  
 C0807  
 C0809  
 C0811  
 C0813  
 C0815  
 C0817  
 C0819  
 C0821  
 C0823  
 C0825  
 C0827  
 C0829  
 C0831  
 C0833  
 C0835  
 C0837  
 C0839  
 C0841  
 C0843  
 C0845  
 C0847  
 C0849  
 C0851  
 C0853  
 C0855  
 C0857  
 C0859  
 C0861  
 C0863  
 C0865  
 C0867  
 C0869  
 C0871  
 C0873  
 C0875  
 C0877  
 C0879  
 C0881  
 C0883  
 C0885  
 C0887  
 C0889  
 C0891  
 C0893  
 C0895  
 C0897  
 C0899  
 C0901  
 C0903  
 C0905  
 C0907  
 C0909  
 C0911  
 C0913  
 C0915  
 C0917  
 C0919  
 C0921  
 C0923  
 C0925  
 C0927  
 C0929  
 C0931  
 C0933  
 C0935  
 C0937  
 C0939  
 C0941  
 C0943  
 C0945  
 C0947  
 C0949  
 C0951  
 C0953  
 C0955  
 C0957  
 C0959  
 C0961  
 C0963  
 C0965  
 C0967  
 C0969  
 C0971  
 C0973  
 C0975  
 C0977  
 C0979  
 C0981  
 C0983  
 C0985  
 C0987  
 C0989  
 C0991  
 C0993  
 C0995  
 C0997  
 C0999  
 C1001  
 C1003  
 C1005  
 C1007  
 C1009  
 C1011  
 C1013  
 C1015  
 C1017  
 C1019  
 C1021  
 C1023  
 C1025  
 C1027  
 C1029  
 C1031  
 C1033  
 C1035  
 C1037  
 C1039  
 C1041  
 C1043  
 C1045  
 C1047  
 C1049  
 C1051  
 C1053  
 C1055  
 C1057  
 C1059  
 C1061  
 C1063  
 C1065  
 C1067  
 C1069  
 C1071  
 C1073  
 C1075  
 C1077  
 C1079  
 C1081  
 C1083  
 C1085  
 C1087  
 C1089  
 C1091  
 C1093  
 C1095  
 C1097  
 C1099  
 C1101  
 C1103  
 C1105  
 C1107  
 C1109  
 C1111  
 C1113  
 C1115  
 C1117  
 C1119  
 C1121  
 C1123  
 C1125  
 C1127  
 C1129  
 C1131  
 C1133  
 C1135  
 C1137  
 C1139  
 C1141  
 C1143  
 C1145  
 C1147  
 C1149  
 C1151  
 C1153  
 C1155  
 C1157  
 C1159  
 C1161  
 C1163  
 C1165  
 C1167  
 C1169  
 C1171  
 C1173  
 C1175  
 C1177  
 C1179  
 C1181  
 C1183  
 C1185  
 C1187  
 C1189  
 C1191  
 C1193  
 C1195  
 C1197  
 C1199  
 C1201  
 C1203  
 C1205  
 C1207  
 C1209  
 C1211  
 C1213  
 C1215  
 C1217  
 C1219  
 C1221  
 C1223  
 C1225  
 C1227  
 C1229  
 C1231  
 C1233  
 C1235  
 C1237  
 C1239  
 C1241  
 C1243  
 C1245  
 C1247  
 C1249  
 C1251  
 C1253  
 C1255  
 C1257  
 C1259  
 C1261  
 C1263  
 C1265  
 C1267  
 C1269  
 C1271  
 C1273  
 C1275  
 C1277  
 C1279  
 C1281  
 C1283  
 C1285  
 C1287  
 C1289  
 C1291  
 C1293  
 C1295  
 C1297  
 C1299  
 C1301  
 C1303  
 C1305  
 C1307  
 C1309  
 C1311  
 C1313  
 C1315  
 C1317  
 C1319  
 C1321  
 C1323  
 C1325  
 C1327  
 C1329  
 C1331  
 C1333  
 C1335  
 C1337  
 C1339  
 C1341  
 C1343  
 C1345  
 C1347  
 C1349  
 C1351  
 C1353  
 C1355  
 C1357  
 C1359  
 C1361  
 C1363  
 C1365  
 C1367  
 C1369  
 C1371  
 C1373  
 C1375  
 C1377  
 C1379  
 C1381  
 C1383  
 C1385  
 C1387  
 C1389  
 C1391  
 C1393  
 C1395  
 C1397  
 C1399  
 C1401  
 C1403  
 C1405  
 C1407  
 C1409  
 C1411  
 C1413  
 C1415  
 C1417  
 C1419  
 C1421  
 C1423  
 C1425  
 C1427  
 C1429  
 C1431  
 C1433  
 C1435  
 C1437  
 C1439  
 C1441  
 C1443  
 C1445  
 C1447  
 C1449  
 C1451  
 C1453  
 C1455  
 C1457  
 C1459  
 C1461  
 C1463  
 C1465  
 C1467  
 C1469  
 C1471  
 C1473  
 C1475  
 C1477  
 C1479  
 C1481  
 C1483  
 C1485  
 C1487  
 C1489  
 C1491  
 C1493  
 C1495  
 C1497  
 C1499  
 C1501  
 C1503  
 C1505  
 C1507  
 C1509  
 C1511  
 C1513  
 C1515  
 C1517  
 C1519  
 C1521  
 C1523  
 C1525  
 C1527  
 C1529  
 C1531  
 C1533  
 C1535  
 C1537  
 C1539  
 C1541  
 C1543  
 C1545  
 C1547  
 C1549  
 C1551  
 C1553  
 C1555  
 C1557  
 C1559  
 C1561  
 C1563  
 C1565  
 C1567  
 C1569  
 C1571  
 C1573  
 C1575  
 C1577  
 C1579  
 C1581  
 C1583  
 C1585  
 C1587  
 C1589  
 C1591  
 C1593  
 C1595  
 C1597  
 C1599  
 C1601  
 C1603  
 C1605  
 C1607  
 C1609  
 C1611  
 C1613  
 C1615  
 C1617  
 C1619  
 C1621  
 C1623  
 C1625  
 C1627  
 C1629  
 C1631  
 C1633  
 C1635  
 C1637  
 C1639  
 C1641  
 C1643  
 C1645  
 C1647  
 C1649  
 C1651  
 C1653  
 C1655  
 C1657  
 C1659  
 C1661  
 C1663  
 C1665  
 C1667  
 C1669  
 C1671  
 C1673  
 C1675  
 C1677  
 C1679  
 C1681  
 C1683  
 C1685  
 C1687  
 C1689  
 C1691  
 C1693  
 C1695  
 C1697  
 C1699  
 C1701  
 C1703  
 C1705  
 C1707  
 C1709  
 C1711  
 C1713  
 C1715  
 C1717  
 C1719  
 C1721  
 C1723  
 C1725  
 C1727  
 C1729  
 C1731  
 C1733  
 C1735  
 C1737  
 C1739  
 C1741  
 C1743  
 C1745  
 C1747  
 C1749  
 C1751  
 C1753  
 C1755  
 C1757  
 C1759  
 C1761  
 C1763  
 C1765  
 C1767  
 C1769  
 C1771  
 C1773  
 C1775  
 C1777  
 C1779  
 C1781  
 C1783  
 C1785  
 C1787  
 C1789  
 C1791  
 C1793  
 C1795  
 C1797  
 C1799  
 C1801  
 C1803  
 C1805  
 C1807  
 C1809  
 C1811  
 C1813  
 C1815  
 C1817  
 C1819  
 C1821  
 C1823  
 C1825  
 C1827  
 C1829  
 C1831  
 C1833  
 C1835  
 C1837  
 C1839  
 C1841  
 C1843  
 C1845  
 C1847  
 C1849  
 C1851  
 C1853  
 C1855  
 C1857  
 C1859  
 C1861  
 C1863  
 C1865  
 C1867  
 C1869  
 C1871  
 C1873  
 C1875  
 C1877  
 C1879  
 C1881  
 C1883  
 C1885  
 C1887  
 C1889  
 C1891  
 C1893  
 C1895  
 C1897  
 C1899  
 C1901  
 C1903  
 C1905  
 C1907  
 C1909  
 C1911  
 C1913  
 C1915  
 C1917  
 C1919  
 C1921  
 C1923  
 C1925  
 C1927  
 C1929  
 C1931  
 C1933  
 C1935  
 C1937  
 C1939  
 C1941  
 C1943  
 C1945  
 C1947  
 C1949  
 C1951  
 C1953  
 C1955  
 C1957  
 C1959  
 C1961  
 C1963  
 C1965  
 C1967  
 C1969  
 C1971  
 C1973  
 C1975  
 C1977  
 C1979  
 C1981  
 C1983  
 C1985  
 C1987  
 C1989  
 C1991  
 C1993  
 C1995  
 C1997  
 C1999  
 C2001  
 C2003  
 C2005  
 C2007  
 C2009  
 C2011  
 C2013  
 C2015  
 C2017  
 C2019  
 C2021  
 C2023  
 C2025  
 C2027  
 C2029  
 C2031  
 C2033  
 C2035  
 C2037  
 C2039  
 C2041  
 C2043  
 C2045  
 C2047  
 C2049  
 C2051  
 C2053  
 C2055  
 C2057  
 C2059  
 C2061  
 C2063  
 C2065  
 C2067  
 C2069  
 C2071  
 C2073  
 C2075  
 C2077  
 C2079  
 C2081  
 C2083  
 C2085  
 C2087  
 C2089  
 C2091  
 C2093  
 C2095  
 C2097  
 C2099  
 C2101  
 C2103  
 C2105  
 C2107  
 C2109  
 C2111  
 C2113  
 C2115  
 C2117  
 C2119  
 C2121  
 C2123  
 C2125  
 C2127  
 C2129  
 C2131  
 C2133  
 C2135  
 C2137  
 C2139  
 C2141  
 C2143  
 C2145  
 C2147  
 C2149  
 C2151  
 C2153  
 C2155  
 C2157  
 C2159  
 C2161  
 C2163  
 C2165  
 C2167  
 C2169  
 C2171  
 C2173  
 C2175  
 C2177  
 C2179  
 C2181  
 C2183  
 C2185  
 C2187  
 C2189  
 C2191  
 C2193  
 C2195  
 C2197  
 C2199  
 C2201  
 C2203  
 C2205  
 C2207  
 C2209  
 C2211  
 C2213  
 C2215  
 C2217  
 C2219  
 C2221  
 C2223  
 C2225  
 C2227  
 C2229  
 C2231  
 C2233  
 C2235  
 C2237  
 C2239  
 C2241  
 C2243  
 C2245  
 C2247  
 C2249  
 C2251  
 C2253  
 C2255  
 C2257  
 C2259  
 C2261  
 C2263  
 C2265  
 C2267  
 C2269  
 C2271  
 C2273  
 C2275  
 C2277  
 C2279  
 C2281  
 C2283  
 C2285  
 C2287  
 C2289  
 C2291  
 C2293  
 C2295  
 C2297  
 C2299  
 C2301  
 C2303  
 C2305  
 C2307  
 C2309  
 C2311  
 C2313  
 C2315  
 C2317  
 C2319  
 C2321  
 C2323  
 C2325  
 C2327  
 C2329  
 C2331  
 C2333  
 C2335  
 C2337  
 C2339  
 C2341  
 C2343  
 C2345  
 C2347  
 C2349  
 C2351  
 C2353  
 C2355  
 C2357  
 C2359  
 C2361  
 C2363  
 C2365  
 C2367  
 C2369  
 C2371  
 C2373  
 C2375  
 C2377  
 C2379  
 C2381  
 C2383  
 C2385  
 C2387  
 C2389  
 C2391  
 C2393  
 C2395  
 C2397  
 C2399  
 C2401  
 C2403  
 C2405  
 C2407  
 C2409  
 C2411  
 C2413  
 C2415  
 C2417  
 C2419  
 C2421  
 C2423  
 C2425  
 C2427  
 C2429  
 C2431  
 C2433  
 C2435  
 C2437  
 C2439  
 C2441  
 C2443  
 C2445  
 C2447  
 C2449  
 C2451  
 C2453  
 C2455  
 C2457  
 C2459  
 C2461  
 C2463  
 C2465  
 C2467  
 C2469  
 C2471  
 C2473  
 C2475  
 C2477  
 C2479  
 C2481  
 C2483  
 C2485  
 C2487  
 C2489  
 C2491  
 C2493  
 C2495  
 C2497  
 C2499  
 C2501  
 C2503  
 C2505  
 C2507  
 C2509  
 C2511  
 C2513  
 C2515  
 C2517  
 C2519  
 C2521  
 C2523  
 C2525  
 C2527  
 C2529  
 C2531  
 C2533  
 C2535  
 C2537  
 C2539  
 C2541  
 C2543  
 C2545  
 C2547  
 C2549  
 C2551  
 C2553  
 C2555  
 C2557  
 C2559  
 C2561  
 C2563  
 C2565  
 C2567  
 C2569  
 C2571  
 C2573  
 C2575  
 C2577  
 C2579  
 C2581  
 C2583  
 C2585  
 C2587  
 C2589  
 C2591  
 C2593  
 C2595  
 C2597  
 C2599  
 C2601  
 C2603  
 C2605  
 C2607  
 C2609  
 C2611  
 C2613  
 C2615  
 C2617  
 C2619  
 C2621  
 C2623  
 C2625  
 C2627  
 C2629  
 C2631  
 C2633  
 C2635  
 C2637  
 C2639  
 C2641  
 C2643  
 C2645  
 C2647  
 C2649  
 C2651  
 C2653  
 C2655  
 C2657  
 C2659  
 C2661  
 C2663  
 C2665  
 C2667  
 C2669  
 C2671  
 C2673  
 C2675  
 C2677  
 C2679  
 C2681  
 C2683  
 C2685  
 C2687  
 C2689  
 C2691  
 C2693  
 C2695  
 C2697  
 C2699  
 C2701  
 C2703  
 C2705  
 C2707  
 C2709  
 C2711  
 C2713  
 C2715  
 C2717  
 C2719  
 C2721  
 C2723  
 C2725  
 C2727  
 C2729  
 C2731  
 C2733  
 C2735  
 C2737  
 C2739  
 C2741  
 C2743  
 C2745  
 C2747  
 C2749  
 C2751  
 C2753  
 C2755  
 C2757  
 C2759  
 C2761  
 C2763  
 C2765  
 C2767  
 C2769  
 C2771  
 C2773  
 C2775  
 C2777  
 C2779  
 C2781  
 C2783  
 C2785  
 C2787  
 C2789  
 C2791  
 C2793  
 C2795  
 C2797  
 C2799  
 C2801  
 C2803  
 C2805  
 C2807  
 C2809  
 C2811  
 C2813  
 C2815  
 C2817  
 C2819  
 C2821  
 C2823  
 C2825  
 C2827  
 C2829  
 C2831  
 C2833  
 C2835  
 C2837  
 C2839  
 C2841  
 C2843  
 C2845  
 C2847  
 C2849  
 C2851  
 C2853  
 C2855  
 C2857  
 C2859  
 C2861  
 C2863  
 C2865  
 C2867  
 C2869  
 C2871  
 C2873  
 C2875  
 C2877  
 C2879  
 C2881  
 C2883  
 C2885  
 C2887  
 C2889  
 C2891  
 C2893  
 C2895  
 C2897  
 C2899  
 C2901  
 C2903  
 C2905  
 C2907  
 C2909  
 C2911  
 C2913  
 C2915  
 C2917  
 C2919  
 C2921  
 C2923  
 C2925  
 C2927  
 C2929  
 C2931  
 C2933  
 C2935  
 C2937  
 C2939  
 C29





Figure 1-4: Genetic maps of the bacteriophage-deficient individuals selected in the first four generations of a marker-selected bacteriophage program. The locus for the bacteriophage-deficient phenotype ( $\Delta b$ ) is located on chromosome I.




Figure 1-4: Genetic maps of the leukemia-derived individuals selected in the first four generations of a marker-assisted breeding program. The locus to be mapped (M) is located on chromosome 1.




Figure 2. Recovery of recurrent parent genotype through backcrossing, with or without marker-assisted selection.

Table 1. Proportion and characteristics of plants carrying the genes of interest, in the first four generations of a marker-assisted breeding program.

generation	% chondrichthys	100% identified	no plants	% homozygous recurrent	no heterozygous
------------	-----------------	-----------------	-----------	------------------------	-----------------




Figure 3: Recovery of mutant plants through backcrossing, with or without marker-assisted selection

Table 1: Proportion and characteristics of plants carrying the genes of interest, in the first four generations of a marker-assisted backcross program.

generation	% phenotypically resistant plants	RFLP genotyping				no plants identified <sup>a</sup>	% intermixture content parent genotype				% intermixture chromosome segments <sup>***</sup>			
		RFLP		RFLP			parent genotype		selected parent		mean		selected parent	
		no plants	no plants	no plants	no plants		mean	std dev	std dev	std dev	mean	std dev	std dev	std dev
BC1	40.0	90	61	2840	57	48.72	10.26	10.26	10.26	10.26	11.01	2.17	7.76	0
BC2	44.8	61	22	1943	39	50.52	1.81	1.81	1.81	1.81	10.44	1.44	3.20	3
BC3	44.8	72	16	720	21	50.52	1.81	1.81	1.81	1.81	10.44	1.44	3.20	3
BC4	44.8	55	5	70	25	50.52	0.49	0.49	0.49	0.49	10.44	1.44	3.20	1

<sup>a</sup> Plants for which two or more adjacent markers had missing values were not included in the analysis

<sup>b</sup> Mean value of the five individuals having the first highest percentage of chromosome segments from the parental genotype.

<sup>c</sup> Including the segment carrying the transgene construct.

comprising the *Rt* locus. It also displays 99.36% of homozygous recurrent-parent-genotype. The remaining 0.64% corresponds to the average relative length of the chromosome segment containing the *Rt* locus, which depends on the two flanking markers chosen.

The mean percentage of homozygous recurrent-parent-genotype of the BC<sub>1</sub> generation was slightly lower than the expected 50%. This can be explained by linkage drag around the *Rt* locus, given that this percentage was computed based only on plants selected for heterozygosity at the *Rt* locus. For all other backcross generations the mean percentage of homozygous recurrent-parent-genotype was much higher than what would have been observed, should no selection have been done (Figure 2).

The percentage of homozygous recurrent-parent-genotype of the selected plant (Table 1) and the average of the five largest values (Table 1) were always very similar to one another, and much superior to the population mean value (Figure 2). The percentage of homozygous recurrent-parent-genotype of the selected plant was found only once, in the BC<sub>2</sub> generation, to be smaller than the average of the five largest values. This corresponded to the only time when the selected plant was not the one with the maximum percentage of homozygous recurrent-parent-genotype. The plant had been selected because it displayed a favorable recombination on one side of the *Rt* locus (Figure 1).

The percentage of homozygous recurrent-parent-genotype of the selected BC<sub>1</sub> plant was almost equal to that of an unselected BC<sub>2</sub>, that of the selected BC<sub>2</sub> was larger than that of an unselected BC<sub>2</sub>, that of the selected BC<sub>3</sub> was barely smaller than that of an unselected BC<sub>4</sub>, and that of the selected BC<sub>4</sub> was equal to that of the "perfect" backcross-derived plant, given the set of markers that was used. Such rates of recurrent parent genotype recovery are consistent with results of simulation analyses. Jarosz *et al.* (1994) who used the maize genome as a model reported that three backcross generations and 80 markers were needed to recover 99% of recurrent parent genotype.

#### Number of donor chromosome segments

The number of heterozygous chromosomal segments decreased from one backcross generation to the next. Plants selected at each generation were not necessarily those which had the lowest number of heterozygous chromosomal segments (Table 1). However, with the set of markers used, BC<sub>3</sub> and BC<sub>4</sub> plants were recovered which contained only one heterozygous chromosomal segment: that comprising the *Rt* locus.

#### Linkage drag

Linkage drag around the *Rt* locus was estimated, relative to the length of chromosome 1. Its value was found to lie between 24.8 and 48.4% for the selected BC<sub>1</sub> individual, between 17.6 and 34.8% for the selected BC<sub>2</sub>, between 7.0 and 24.0% for the selected BC<sub>3</sub>, and between 0.0 and 4.4% (respectively 0.0 and 14.5 cM) for the selected BC<sub>4</sub>.

The two values given for each *g* correspond to extreme positions of flanking the transgene construct locus BC<sub>4</sub> is likely to be less than 1.5% appear to be somewhat high, reflect drag, it is much lower than what (Stam and Zeven 1981; Tankley *et al.* of tomato cultivars obtained by a Tankley (1989) found that the sizes cM.

#### Conclusion

These results clearly demonstrate quality advantages over classical through backcrossing. Only four to five years and a half from plant genotypically fully converted. New genotype could protect over vast areas appropriate protocols and resources allocated.

Comparison of BC<sub>4</sub>-derived 1 markers and agronomic performance order to confirm the completeness to

#### References

- ALLARD, R.W. (1986) Principles of plant breeding. Academic Press, New York.
- HALLAUER, A.R., and J.B. MIRANDA. (1989) University Press, Ames, IA.
- HOSPITAL, F., C. COMVALENT, and F.J. GARCIA. (1992) Genetica 132:1199-1210.
- JAROSZ, J.G., W. DEBRAVIS, and S.J.O. (1994) Select backcross programs by computer on the plant genome. Schering International.
- KOZIEL, M.G., G.L. INELAND, C. PO. RAWSON, R. OSEAR, M. WILLIAMS, SUPERVISION, M.R. MECHER, S. M. IVOLA (1993) Field performance of derived from *Radiata* divergent. *St*
- MURRAY, M.G., Y.M.A., J. ROMERO. (1994) Segment length polymorphism what

homologous recurrent-parent-genotype. The relative length of the chromosome between the two flanking markers chosen.

The parent-genotype of the BC<sub>1</sub> generation was explained by linkage drag around the selected gene based only on plants selected for one generation; the mean percentage of linkage drag was higher than what would have been expected.

The parent-genotype of the selected plant (Table 1) were always very similar to the parent value (Figure 2). The percentage of linkage drag of the selected plant was found only once, in the five largest values. This corresponded to one with the maximum percentage of linkage drag selected because it displayed a linkage drag value.

The parent-genotype of the selected BC<sub>1</sub> plant of the selected BC<sub>2</sub> was larger than that of the "perfect" backcross-derived plant. The rate of recurrent parent genotype was lower. Jarboe *et al.* (1994) who used backcross generations and 80 markers found linkage drag.

Linkage drag decreased from one backcross generation to the next. The linkage drag was not necessarily those which were selected (Table 1). However, with linkage drag which contained only one gene at the locus.

Linkage drag relative to the length of chromosome was 1.3% for the selected BC<sub>1</sub> individual, 2.0% and 24.0% for the selected BC<sub>2</sub> and BC<sub>3</sub> respectively.

The two values given for each generation are extreme values of linkage drag, which correspond to extreme positions of the crossing-over in the marker-defined intervals flanking the transgene construct locus. Therefore the true linkage drag value of the selected BC<sub>2</sub> is likely to be less than 1.3% of the genome. Although this maximum value may appear to be somewhat high, reflecting the limited selection pressure put here on linkage drag, it is much lower than what would be expected from classical backcross programs (Shan and Zeev 1981; Tanksley *et al.* 1989). Practically, in a study of *Tm-2* conversions of tomato cultivars obtained by a large number of classical backcross cycles, Young and Tanksley (1989) found that the sizes of the introgressed fragments ranged between 4 and 51 cM.

## Conclusion

These results clearly demonstrate that molecular markers provide important time and quality advantages over classical procedures for the production of near-isogenic lines through backcrossing. Only four backcross generations were necessary to recover, in less than a year and a half from planting of the BC<sub>1</sub>'s, individuals which appeared to be genetically fully converted. Nevertheless, it is likely that recovery of recurrent parent genotype could proceed even faster than in the experiment described herein, should the appropriate protocol and resources (population size, number and position of markers) be allocated.

Comparison of BC<sub>2</sub>-derived lines with the recurrent parent for both morphological markers and agronomic performance (including hybrid performance) will be performed in order to confirm the completeness of the conversion.

## References

- ALLARD, R.W. (1968) *Principles of plant breeding*. Wiley, New York, NY.
- HALLAUER, A.R., and J.R. BRAND, Jr. (1961) *Quantitative genetics in maize breeding*. Iowa State University Press, Ames, IA.
- HOSPITAL, F., C. CHEVALE, and P. MULSANT (1992) Using markers in gene introgression breeding programs. *Genetics* 122:1199-1210.
- JARBOE, S.G., W.D. BEAVIS, and S.J. OPENSHAW (1994) Prediction of responses to selection in marker-assisted backcross programs by computer simulation. In: *Abstracts of the second international conference on the plant genome*. Science International Inc. 34.
- KOZEL, M.G., G.J. BELAND, M.R. BOWMAN, M.R. CAROZZE, R. CRISHAW, L. CROSSLAND, I. DAWSON, N. DEBAL, M. FELL, S. KADWELL, K. LAURIS, E. LEWIS, D. MADDOX, K. McPHERSON, M.R. MOORE, E. MUELLER, R. RHOODES, G.W. WARREN, M. WRIGHT, and S.V. SYOLA (1993) Field performance of elite transgenic maize plants expressing an insecticidal protein derived from *Bacillus thuringiensis*. *Biotechnology* 11:194-200.
- MURRAY, M.G., Y.M.A. J. MONTEIRO-SILVEIRA, D.P. WEST, and J.M. CRAMER (1988) Recombination segment length polymorphism: what are they and how can breeders use them? In: D. W. HILLIS *et al.*



Proceedings of the 43rd annual corn and sorghum industry research conference. American Seed Trade Association 43/72-87.

STAM, P., and C. ZEVEN (1981) The domestic population of the donor genome in near-isogenic lines of self-fertile lines bred by backcrossing. *Biometrics* 36:223-238.

TANLEY, S.D., N.D. YOUNG, A.H. PATTERSON, and M.W. SONNERSALE (1989) RFLP mapping in plant breeding: new tools for an old science. *BioTechnology* 7:257-264.

YOUNG, N.D., and S.D. TANLEY (1989) RFLP analysis of the site of chromosome segments retained around the 7m-2 locus of corn during backcross breeding. *Theor. Appl. Genet.* 77:353-359.

C

# Marker-assisted Selection in Backcross Breeding

S.J. Openshaw

Pioneer Hi-Bred Intl. Inc., P.O. Box 1004, Johnston, IA 50131

S.G. Jarboe<sup>1</sup>

CIMMYT, Lisboa 27, Apdo. Postal 6-641, 06600 Mexico, D.F., Mexico

W.D. Beavis

Pioneer Hi-Bred Intl. Inc., P.O. Box 1004, Johnston, IA 50131

Notice: This material may be protected by copyright law (Title 17, U.S. Code)

**Abstract.** The backcross breeding procedure has been used widely to transfer simply inherited traits into elite genotypes. Genetic markers can increase the effectiveness of backcrossing by 1) increasing the probability of obtaining a suitable conversion, and 2) decreasing the time required to achieve an acceptable recovery. Simulation and field results indicated that, for a genome consisting of ten 200-cM chromosomes, basing selection on 40 or 80 markers in 50 BC individuals that carry the allele being transferred can reduce the number of backcross generations from about seven to three.

The backcross breeding procedure has been used widely to transfer simply inherited traits into elite genotypes. Usually, the trait being transferred is controlled by a single gene, but highly heritable traits that are more complexly inherited have also been transferred successfully by backcrossing; for example, maturity in maize (Johns and Sauer, 1961; Shaver, 1976). Today, backcrossing is being used to transfer genes introduced by such techniques as transformation or mutation into appropriate germplasm.

Several plant breeding textbooks give good descriptions of the backcross procedure (Allard, 1960; Fehr, 1977). A donor parent (DP) carrying a trait of interest is crossed to the recurrent parent (RP), an elite line that is lacking the trait. The F<sub>1</sub> is crossed back to the RP to produce the BC<sub>1</sub> generation. In the BC<sub>1</sub>, and subsequent backcross generations, selected individuals carrying the gene being transferred are backcrossed to the RP. The expected proportion of DP genome is reduced by half with each generation of backcrossing. Ignoring effects of linkage to the selected DP allele being transferred, the percentage recurrent parent (%RP) genome expected in each backcross generation is calculated as:

$$\%RP = 100 [1 - (0.5)^n]$$

where  $n$  is the number of backcrosses.

Backcrossing of selected plants to the RP can be repeated each cycle until a line is obtained that is essentially a version of the RP that includes the introgressed allele. After six backcrosses, the expected recovery is >96% (Table 1).

Until recently, discussions of the recovery of the RP genome during backcrossing have emphasized the expected values for

%RP shown in Table 1, and have largely ignored the genetic variation for %RP that exists around the expected mean. With the development of genetic markers capable of providing good genome coverage, there has been interest in taking advantage of that variation to increase the efficiency of backcrossing.

Selection for RP marker alleles can increase greatly the effectiveness of backcross programs by allowing the breeder to: 1) select backcross plants that have a higher proportion of RP genome, and 2) select backcross individuals that are better converters of a mapped donor allele being transferred (i.e., select for less linkage drag). Expressed in practical terms, using genetic markers to assist backcrossing can 1) increase the probability of obtaining a suitable conversion, and 2) decrease the time required to achieve an acceptable recovery.

Issues to consider when planning a marker-assisted backcross program include 1) the time advantage of using markers to assist backcrossing, 2) the number of markers needed, and 3) the number of genotypes to evaluate. In this report, we use results from previous literature, computer simulation, and empirical studies to provide some guidelines.

Table 1. Expected recovery of recurrent parent (RP) genome during backcrossing, assuming no linkage to the gene being transferred.

Generation	%RP
F <sub>1</sub>	50.0000
BC <sub>1</sub>	75.0000
BC <sub>2</sub>	87.5000
BC <sub>3</sub>	93.7500
BC <sub>4</sub>	96.8750
BC <sub>5</sub>	98.4375
BC <sub>6</sub>	99.2188
BC <sub>7</sub>	99.6094

<sup>1</sup>Presently with Purdue University, West Lafayette, Ind.

## Analysis of Molecular Marker Data

## Materials and methods

The maize genome was the model for the simulation. The simulated genome consisted of 200-cM chromosomes. Simulation of crossing over was based on a Poisson distribution with a mean of 2.0 ( $\lambda = 2$ ) (Hasson, 1959), which, on average, generated one cross over for every 100-cM length. The simulations reported here assume no interference. Codominant genetic markers were evenly distributed in the genome and sites of the donor genes were randomly assigned to genome locations. Simulations were conducted with the following parameters:

- Number of progeny: 100 or 500.
- Backcross generations:  $BC_1$ ,  $BC_2$ , and  $BC_3$ .
- Number of markers: 20, 40, 80, or 100.
- Number selected to form the next BC generation: 1 or 5.

Selection was based on 1) presence of the donor allele and 2) high %RP. %RP was calculated as the average of the (one or five) selected individuals. Values presented are the mean of 50 simulations.

## Results

In the computer simulation study, all methods modeled greatly increased the speed of recovering the RP genome compared to the expected recovery with no marker-assisted selection (compare Tables 1 and 2). At least 80 markers were required to recover 99% of the RP genome in just three BC generations (Table 2). The use of at least 80 markers and 500 progeny allowed recovery of 98% RP in just two BC generations. Response to selection was diminished only slightly by spreading the effort over five selections. Using markers, the number of backcross generations needed to convert an inbred is

reduced from about seven to three.

By the  $BC_3$  generation, there appears to be no practical advantage to using 500 vs. 100 individuals. If the presence of the donor trait in the backcross individuals can be ascertained before markers are genotyped, then only half the number of individuals indicated in the tables will need to be analyzed.

When a small number of markers are used, they quickly become non-informative; i.e., selection causes the marker loci to become fixed for the RP type before the rest of the genome is fully converted (Table 3; Hospital et al., 1992). This situation was most prominent in the larger populations, where a higher selection intensity placed more selection pressure upon the marker loci. Accordingly, it is of interest to consider how closely the estimation of %RP based on markers reflects the actual genome composition. The combination of estimation of %RP based on fewer markers and subsequent selection tends to bias the estimates upward (compare Tables 2 and 3).

The results from the simulation compare well with real field data. In a typical example, 50  $BC_3$  plants carrying the gene being transferred were genotyped at 83 polymorphic RFLP loci (note that this corresponds to a population size of 100 unselected plants in Tables 2 and 3). The five best  $BC_3$  recoveries had estimated %RP values of 83.9%, 82.7%, 82.0%, 81.4%, and 81.2%. After evaluating 10  $BC_3$  plants from each selected  $BC_3$ , the best  $BC_3$  recovery had an estimated %RP of 94.6%.

## Discussion

The simulations (Table 2; Hospital et al., 1992) and our experience indicate that four markers per 200-cM chromosome is adequate to greatly increase the effectiveness of selection in the  $BC_3$ . However, using only four markers per 200 cM will likely make it very difficult to map the location of the gene of interest. Adequate summation of this data is an important

Table 2. Percent recurrent parent genome during marker-assisted backcrossing.

Generation	100 Progeny No. markers				500 Progeny No. markers			
	20	40	80	100	20	40	80	100
<i>One selected</i>								
$BC_1$	64.3	64.3	64.3	88.0	89.9	90.7	90.3	90.3
$BC_2$	93.0	93.3	95.4	97.3	96.5	97.7	98.5	98.6
$BC_3$	97.4	97.6	98.9	99.3	97.7	98.3	99.4	99.3
<i>Five selected</i>								
$BC_1$	82.9	83.1	84.9	84.7	87.7	88.1	88.9	88.9
$BC_2$	93.7	93.0	93.8	93.7	95.5	96.8	97.8	97.9
$BC_3$	97.1	98.3	98.8	98.9	97.3	98.3	99.3	99.3

Table 3. Estimates of percent recurrent parent genome, based on marker loci.

Generation	100 Progeny No. markers				500 Progeny No. markers			
	20	40	80	100	20	40	80	100
<i>One selected</i>								
$BC_1$	98.7	97.4	95.6	97.2	100.0	99.1	98.6	98.0
$BC_2$	100.0	99.8	99.3	99.5	100.0	100.0	99.9	98.2
<i>Five selected</i>								
$BC_1$	96.4	96.5	96.2	95.8	100.0	99.5	98.3	98.2
$BC_2$	99.9	99.8	99.3	99.1	100.0	100.0	99.9	99.8

part of a marker-assisted backcross program. Ideally, the markers used can supply data that can be represented as alleles of loci with known map position. Estimation of WRF, mapping the position of the locus of interest, and graphical display of the results (Young and Tanksley, 1989) are all useful in understanding and controlling the specific backcross experiment being conducted.

It appears that, with the use of genetic markers, the portion of the RP genome that is not linked to the allele being transferred can be recovered quickly and with confidence. The recovery of RP will be slower on the chromosome carrying the gene of interest. A considerable amount of linkage drag is expected to accompany selection for the DF allele in a backcross program. For a locus located in the middle of a 200-cM chromosome, the length of the DF chromosome segment accompanying selection is expected to be 124, 63, and 21 cM in the BC<sub>1</sub>, BC<sub>2</sub>, and BC<sub>3</sub> generations, respectively (Hanson, 1959; Naveira and Barbudillo, 1992). Our observations support the recommendation of Hospital et al. (1992) that preference be given to the selection for recombinants proximal to the allele of interest, but that selection for recovery of the RP elsewhere in the genome also be considered. This two-stage selection can probably be done quite effectively and has by the breeder once the data is adequately summarized; however, Hospital et al.

suggest ways to incorporate the two criteria into a selection index such that each component of selection is assured appropriate weighting.

Use of genetic markers can greatly increase the effectiveness of backcrossing, and they should be used in any serious backcrossing program if resources are available to the breeder.

#### Literature Cited

- Allard, R.W. 1960. Principles of plant breeding. Wiley, New York.  
 Fehr, W.F. 1987. Principles of cultivar development: v.1. Theory and technique. Macmillan, New York.  
 Hanson, W.D. 1959. Early generation analysis of length of heterozygous chromosome segments around a locus fixed heterozygous with backcrossing or selfing. *Genetics* 44:843-847.  
 Hospital, F., C. Chevillon, and P. Malouf. 1992. Using markers in gene introgression breeding programs. *Genetics* 132:1199-1210.  
 Ricka, E.H. and J.C. Sentz. 1961. Moving corn-belt germplasm northward. *Ann. Hybrid Corn Industry Conf.* 16:53-58.  
 Shriver, D.L. 1976. Conversions for selfiness in maize inbreds. *Maine Genet. Conf. Nwark*, 50:20-21.  
 Young, N.D. and S.D. Tanksley. 1989. Restriction fragment length polymorphism maps and the concept of graphical genotypes. *Theor. Appl. Genet.* 77: 95-101.



## Essential Derivation and Dependence

### Practical Information

---

#### WHY THE CONCEPT OF ESSENTIAL DERIVATION?

The 1978 Act of the UPOV Convention (International Union for the Protection of New Varieties of Plants) states that "the authorization by the breeder shall not be required either for the utilization of the [his protected] variety as an initial source of variation for the purpose of creating other varieties or for the marketing of such varieties".

That principle, known as the "breeder's exemption", is essential for continued progress from plant breeding.

However, its implementation has progressively led to some abuses, due to the difficulties involved with assessment of distinctness, based on the text of the Convention (1978) which indicates that, for the basis of a title of protection, "the [new] variety must be clearly distinguishable by one or more important characteristics from any other variety whose existence is a matter of common knowledge ...".

Sometimes, "cosmetic modifications" were enough for protecting a new variety. That was particularly true in the case of mutation of ornamental or fruit plants and of "conversion" by repeated backcrossing of parental lines of hybrid varieties.

In order to improve the situation, in the early 1980's, a debate began on how to improve the system, trying to define "minimum distances" per species, but no consensus was reached. The development of genetic engineering, opened new possibilities for "piracy" of varieties and sped up the revision process of the Convention which, in the Act adopted in 1991, has introduced with the full agreement of breeders' associations, the concept of essential derivation. That concept of essential derivation has two aspects:

- a technical one: the question whether or not a plant variety is to be considered as a variety essentially derived from an initial variety;
- a juridical one: dependence, meaning that no protected acts as defined by the 1991 Act of the UPOV Convention (production, marketing ...) related to the essentially derived variety shall be carried out without the authorization of the owner of the protected initial variety.

#### DEFINITION OF AN ESSENTIALLY DERIVED VARIETY

The 1991 Act of the UPOV Convention states that "a variety shall be deemed to be essentially derived from another variety (the initial variety) when:

- I. It is predominantly derived from the initial variety, or from a variety that is itself predominantly derived from the initial variety, while retaining the expression of the essential characteristics that result from the genotype or combination of genotypes of the initial

variety;

- ii. it is clearly distinguishable from the initial variety and
- iii. except for the differences which result from the act of derivation, it conforms to the initial variety in the expression of the essential characteristics that result from the genotype or combination of genotypes of the initial variety.

Essentially derived varieties may be obtained, for example, by selection of natural or induced mutants, by selection of a somaclonal variant, by selection of variant individual plants in the initial variety, by backcrossing or by transformation (genetic engineering).

ASSINSEL interprets the definition given in the Convention as follows:

a) The technical aspects (matter of facts)

For a variety to be considered as essentially derived, it must fulfil three requirements in relation to the initial variety while retaining the expression of the essential characteristics of the initial variety:

- i. clear distinctness in the sense of the UPOV Convention
- ii. conformity to the initial variety in the expression of the essential characteristics that result from the genotype or combination of genotypes of the initial variety
- iii. predominant derivation from an initial variety.

If one of these requirements is not fulfilled, there is no essential derivation.

The methods of breeding that can be regarded as leading to an essentially derived variety (see the above-mentioned explanatory list) may differ from species to species or even within a species. This may result in different thresholds being required to characterize essential derivation. Thus, conformity should be judged on a species-by-species or even within a species basis.

b) The juridical aspect

The principle of dependence only exists in favour of a protected variety. This means that:

- i. the initial variety must be a protected one
- ii. dependence can only exist from one protected variety alone
- iii. an essentially derived variety can be directly derived from the initial variety or from a variety that is itself predominantly derived from the initial variety. It is possible to have a "cascade" of derivation. However, each essentially derived variety shall only be dependent on one, the protected initial variety. A cascade of dependence shall not exist, the principle having been introduced to better protect the breeder of the initial variety and not those having made derivations from his work.

## ASSESSMENT OF ESSENTIAL DERIVATION

The assessment of essential derivation needs to take into account the three criteria mentioned above:

- ⊕ clear distinctness in the sense of the UPOV Convention
- ⊕ conformity to the initial variety in the expression of the essential characteristics that result from the genotype or the combination of genotypes of the initial variety
- ⊕ predominant derivation from an initial variety.

The first criterion will be decided upon by the office in charge of granting a right to the breeder of the variety, according to the UPOV rule of distinctness.

The second criterion could be based on reliable phenotypic characteristics and/or on reliable molecular characteristics: either close relationship in general which could lead to a "conformity threshold" parallel to the minimum distance threshold used for distinctness or only small differences in some simply inherited characteristics. If this second criterion is considered as fulfilled, then, we have to assess the third one, which is "predominant derivation from an initial variety".

The third criterion, predominant derivation from an initial variety, implies that the initial variety or products essentially derived therefrom have been used in the breeding process.

In order to prove that use, various criteria or a combination thereof may be used:

- ⊕ combining ability
- ⊕ phenotypic characteristics
- ⊕ molecular characteristics.


These criteria will have to be handled differently from their use for assessment of distinctness. Whatever solution retained, one will probably have to use distance coefficients to define thresholds. Up to now, ASSINSEL has essentially worked on thresholds based on distances measured by molecular markers. Geneticists and statisticians consider that technically it is equally possible to measure distance coefficients using phenotypic markers. However, the process would probably be more difficult due to environmental factors and much more expensive: necessity of several testing locations during several years. However, if breeders prefer to use morphological markers instead of molecular markers, that should be possible.

The interest of using combining ability and the heterosis level will strongly depend on the crop. Thresholds will also be necessary.

The various ASSINSEL Sections are considering the establishment of thresholds for characterization of essential derivation according to the following general principle:

- ⊕ One should propose, species by species, a first threshold below which a variety should be considered as non-essentially derived from an initial variety and a second threshold of conformity above which the new variety should be considered as essentially derived, except if the breeder can prove, by clear evidence, that he has started from independent germplasm.
- ⊕ Between those two thresholds, the derivation could be disputable and the breeder of the putative essentially derived variety should have to give, in case of amicable negotiation or arbitration, information on the origin of the new variety. Should that information be unsatisfactory, the tribunal or of arbitrators/conciliators agreed on by both parties may request breeding records be provided for their examination.

This approach may be diagrammed as follows:



Some breeders are developing such scheme and call the zone No.1 "green zone", in which breeders would have freedom to operate. Zone No.3, the "red zone", where the breeder would know, according to his breeding materials, if his new variety is obviously essentially derived and dependent. Zone No.2 is where there would be uncertainty and where discussion may be appropriate. The threshold levels would be established first as an experiment. They could be further modified according to the experience acquired in the implementation of the scheme.

While this approach may be worthwhile, it also presents some obvious difficulties:

- Breeders have so far been unable to agree on threshold levels for any species;
- Even if the thresholds adopted by the industry had merit, they will not represent an absolute certainty and a court of law could pass judgment on other bases or guidelines.

Nevertheless, this approach does provide some framework in which breeders might proceed.

#### CONSEQUENCES FOR THE BREEDERS

The concepts of derivation and dependence do not, fortunately, abolish the "breeder's exemption" which is still stated in the 1991 Act. However, "cosmetic" improvement or plagiarism, which could sometimes have allowed the creation of distinct varieties in the sense of the UPOV Convention, will no longer allow the creation of independent varieties. The consequences for the breeders, the farmers and biological diversity more broadly should be positive and will certainly impact the breeder's work.

##### a) Choice of the parents

Breeders should be certain of their legal access and freedom to use all parent materials employed in their breeding programs. They would have to pay more attention to the results of their breeding work when working with protected varieties within the "breeder's exemption".

##### b) Breeding methods

Any conventional breeding method could, in theory, provide an essentially derived



## Essential Derivation and Dependence - Practical Information

variety. Certain methods appear to give a higher risk of developing essentially derived varieties. Among these methods we include:

- ④ natural or induced mutations;
- ④ repeated backcrosses; (discussions still continue on the number of backcrosses which could lead to an essentially derived variety. As shown in the French text of the 1991 Convention, which is of evidence, the authors of the Convention had in mind at least two backcrosses, the word being written in plural. However, it must be noted that the selection pressure exerted after the backcross(es) can have an important effect on the final result).
- ④ selection in an existing variety, for example the choice of clones in a synthetic variety;
- ④ transformation by genetic engineering.

c) Development of technical information

Conformity thresholds for essential derivation, such as presented above, can be defined in the frame of professional agreement (which would be the solution) or, in a case-by-case basis, in decisions by courts of law. In either case, thresholds will come to exist in the years ahead. To know their freedom to operate in relation to such thresholds, breeders will need:

- ④ a good knowledge of the range of phenotypic, molecular and physiological variability of varieties present in the market;
- ④ to know the phenotypic, molecular and physiological profiles of their genetic material and their experimental varieties, as well as their breeding histories and documentation of legal access.

Breeders will need to employ the tools necessary for assessing such profiles in their research programs. Such tools will not only be used for the protection of intellectual property, but should also promote improvement of breeding efficiency.

d) Keeping of breeding books

Conformity thresholds only, at least in the zone of uncertainty (orange zone), will not allow a decision on derivation and dependence. In case of litigation, information on parental material and breeding methods will be needed. Thus, breeders will need to maintain clear and accurate breeding records. We encourage breeders to seek competent professional legal advice on the best ways to develop and maintain these important records.

### What is an "Essentially Derived Variety"?

The concept of essentially derived variety was introduced into the 1991 Act of the UPOV Convention in order to avoid plagiarism through mutation, multiple back-crossing and to fill the gap between Plant Breeder's Rights and patents, gap which was becoming important due to the development of the use of patented genetic traits in genetic engineering.

An essentially derived variety is a variety which is distinct and predominantly derived from a protected initial variety, while retaining the essential characteristics of that initial variety.

As indicated as an example in the UPOV Convention, essentially derived varieties may be obtained by the selection of a natural or induced mutant, or of a somaclonal variant, the selection of a variant individual from plants of the initial variety, back-crossing, or transformation by genetic engineering.

The commercialization of an essentially derived variety needs the authorization of the owner of the rights vested in the initial variety.

The concept of essentially derived variety does not at all abolish the Breeder's Exemption, as free access to protected plant varieties for breeding purposes is maintained. It is not a threat to biodiversity. On the contrary, it favors biodiversity, encouraging breeders developing and marketing original varieties.